

ANAPHYLAXIS POLICY



PURPOSE

To explain to Frankston Heights Primary School parents, carers, staff and students the processes and procedures in place to support students diagnosed as being at risk of suffering from anaphylaxis. This policy also ensures that Frankston Heights Primary School is compliant with Ministerial Order 706 and the Department's guidelines for anaphylaxis management.

SCOPE

This policy applies to:

- all staff, including casual relief staff, canteen operators and volunteers
- all students who have been diagnosed with anaphylaxis, or who may require emergency treatment for an anaphylactic reaction, and their parents and carers.

POLICY

School Statement

Frankston Heights Primary School will fully comply with Ministerial Order 706 and the associated guidelines published by the Department of Education and Training.

Anaphylaxis

Anaphylaxis is a severe allergic reaction that occurs after exposure to an allergen. The most common allergens for school-aged children are nuts, eggs, cow's milk, fish, shellfish, wheat, soy, sesame, latex, certain insect stings and medication.

Symptoms

Signs and symptoms of a mild to moderate allergic reaction can include:

- swelling of the lips, face and eyes
- hives or welts
- tingling in the mouth.

Signs and symptoms of anaphylaxis, a severe allergic reaction, can include:

- difficult/noisy breathing
- swelling of tongue
- difficulty talking and/or hoarse voice
- wheeze or persistent cough
- persistent dizziness or collapse
- student appears pale or floppy
- abdominal pain and/or vomiting.

Symptoms usually develop within ten minutes and up to two hours after exposure to an allergen, but can appear within a few minutes.

Treatment

Adrenaline given as an injection into the muscle of the outer mid-thigh or inhaled through the nose is the first aid treatment for anaphylaxis.

Individuals diagnosed as being at risk of anaphylaxis are prescribed an adrenaline autoinjector for use in an emergency. These adrenaline autoinjectors are designed so that anyone can use them in an emergency.

Individual Anaphylaxis Management Plans

All students at Frankston Heights Primary School who are diagnosed by a medical practitioner as being at risk of suffering from an anaphylactic reaction must have an Individual Anaphylaxis Management Plan. When notified of an anaphylaxis diagnosis, the FHPS First Aid Officer is responsible for developing a plan in consultation with the student's parents/carers.

Where necessary, an Individual Anaphylaxis Management Plan will be in place as soon as practicable after a student enrolls at Frankston Heights Primary School and where possible, before the student's first day.

Parents and carers must:

- obtain an ASCIA Action Plan for Anaphylaxis (RED) from the student's medical practitioner and provide a copy to the school as soon as practicable
- immediately inform the school in writing if there is a relevant change in the student's medical condition and obtain an updated ASCIA Action Plan for Anaphylaxis (RED)
- provide an up-to-date photo of the student for the ASCIA Action Plan for Anaphylaxis (RED) and provide it to the school each time it is updated
- provide the school with a current adrenaline device for the student that has not expired
- participate in annual reviews of the student's Individual Anaphylaxis Management Plan that is prepared by the school.

Each student's Individual Anaphylaxis Management Plan must include:

- information about the student's medical condition that relates to allergies and the potential for anaphylactic reaction, including the type of allergies the student has
- information about the signs or symptoms the student might exhibit in the event of an allergic reaction based on a written diagnosis from a medical practitioner
- strategies to minimise the risk of exposure to known allergens while the student is under the care or supervision of school staff, including in the school yard, at camps and excursions, or at special events conducted, organised or attended by the school
- the name of the person(s) responsible for implementing the risk minimisation strategies, which have been identified in the Individual Anaphylaxis Management Plan
- information about where the student's medication is stored
- the student's emergency contact details
- an up-to-date ASCIA Action Plan for Anaphylaxis (RED) completed by the student's medical practitioner.

Review and updates to Individual Anaphylaxis Management Plans

A student's Individual Anaphylaxis Management Plan will be reviewed and updated on an annual basis in consultation with the student's parents/carers. The plan will also be reviewed and, where necessary, updated in the following circumstances:

- as soon as practicable after the student has an anaphylactic reaction at school
- if the student's medical condition, insofar as it relates to allergy and the potential for anaphylactic reaction, changes

- when the student is participating in an off-site activity, including camps and excursions, or at special events including fairs and concerts.

Our school may also consider updating a student's Individual Anaphylaxis Management Plan if there is an identified and significant increase in the student's potential risk of exposure to allergens at school.

Location of Plans and Adrenaline Autoinjectors

A copy of each student's Individual Anaphylaxis Management Plan will be stored with their ASCIA Action Plan for Anaphylaxis (RED) in the First Aid room, together with the student's adrenaline device. Adrenaline devices must be labelled with the student's name.

Risk Minimisation Strategies

To reduce the risk of a student suffering from an anaphylactic reaction at Frankston Heights Primary School we have put in place the following strategies:

- staff and students are regularly reminded to wash their hands after eating
- students are discouraged from sharing food (reminders published in school newsletter)
- rubbish bins at school are to remain covered with lids to reduce the risk of attracting insects
- gloves can be worn or tongs used when picking up papers or rubbish in the playground
- hands are to be thoroughly washed following a yard clean up
- school canteen staff are trained in appropriate food handling to reduce the risk of cross-contamination
- home group classes will be informed of allergens specific to students in their class that must be avoided in advance of class parties, events or birthdays
- a general use adrenaline autoinjector will be stored in the First Aid room
- planning for off-site activities will include risk minimisation strategies for students at risk of anaphylaxis including supervision requirements, appropriate number of trained staff, emergency response procedures and other risk controls appropriate to the activity and students attending.

Adrenaline Autoinjectors for General Use

Frankston Heights Primary School will maintain a supply of adrenaline autoinjector(s) for general use, as a back-up to those provided by parents and carers for specific students, and also for students who may suffer from a first time reaction at school.

There are currently 4 adrenaline devices approved by the Therapeutic Goods Administration for use in Australia: the EpiPen®, the Anapen®, Jext® and Neffy®. All devices can be used when provided by families for students, however, the principal or allocated staff member can only use EpiPen®, Anapen® or Jext® adrenaline autoinjector for general use. For more information about which autoinjector to purchase for general use, refer to [Adrenaline autoinjectors for general use](#).

Adrenaline autoinjectors for general use will be stored in the First Aid room and labelled "general use".

The First Aid Officer is responsible for arranging the purchase of adrenaline autoinjectors for general use, and will consider:

- the number of students enrolled at Frankston Heights Primary School at risk of anaphylaxis
- the accessibility of adrenaline autoinjectors supplied by parents/carers
- the availability of a sufficient supply of autoinjectors for general use in different locations at the school, as well as at camps, excursions and events
- the limited life span of adrenaline autoinjectors, and the need for general use adrenaline autoinjectors to be replaced when used or prior to expiry

- the weight of the students at risk of anaphylaxis to determine the correct dosage of adrenaline autoinjector/s to purchase.

Emergency Response

In the event of an anaphylactic reaction, the emergency response procedures in this policy must be followed, together with the school's general first aid procedures, emergency response procedures and the student's Individual Anaphylaxis Management Plan.

A complete and up-to-date list of students identified as being at risk of anaphylaxis is maintained by the First Aid Officer and stored in the First Aid room. For camps, excursions and special events, a designated staff member will be responsible for maintaining a list of students at risk of anaphylaxis attending the special event, together with their Individual Anaphylaxis Management Plans and adrenaline autoinjectors, where appropriate.

If a student experiences an anaphylactic reaction at school or during a school activity, school staff must:

Step	Action
1.	• Lay the person flat. • Do not allow them to stand or walk. • If breathing is difficult, allow them to sit. • Be calm and reassuring. • Do not leave them alone. • Seek assistance from another staff member or reliable student to locate the student's adrenaline autoinjector or the school's general use autoinjector, and the student's Individual Anaphylaxis Management Plan, stored in the First Aid Room. • If the student's plan is not immediately available, or they appear to be experiencing a first-time reaction, continue with steps 2 to 5 and refer to the ASCIA First Aid Plan for Anaphylaxis (Orange), stored with the school's adrenaline autoinjector for general use in the First Aid Room, if immediately accessible.
2.	Administer an EpiPen or EpiPen Jr • Remove the EpiPen from its plastic container. • Form a fist around the EpiPen and pull off the blue safety release (cap). • Hold the student's leg still and place the orange end against the student's outer mid-thigh (with or without clothing). • Push down firmly until a click is heard or felt, and hold in place for 3 seconds. • Remove the EpiPen. • Note the time the EpiPen was administered. • Retain the used EpiPen and hand it to the ambulance paramedics, together with the time of administration. OR Administer an Anapen® 500 • Pull off the black needle shield. • Pull off the grey safety cap (from the red button). • Place the needle end firmly against the student's outer mid-thigh at a 90-degree angle (with or without clothing). • Press the red button until it clicks and hold for 3 seconds. • Remove the Anapen®. • Note the time the Anapen was administered. • Retain the used Anapen and hand it to the ambulance paramedics, together with the time of administration. OR

	Administer Jext 150 or 300 • Form fist around Jext and pull off yellow cap. • Place black injector tip against outer-mid thigh (with or without clothing). • Push black tip firmly until a click is heard and hold in place for 3 seconds. • Remove Jext. • Note the time the Jext device is administered. • The used adrenaline device must be handed to the ambulance paramedics along with the time of administration. OR Administer Neffy® 1mg or 2mg • Hold the nasal spray with your thumb on the bottom of the plunger and a finger on either side of the nozzle. • Do not pull or push on the plunger. Do not test or prime (pre-spray). Each Neffy nasal spray contains only one spray. • Place the nozzle of the nasal spray into a nostril until your fingers touch the nose. • For smaller nostrils, aim for your fingers to touch the nose. • Keep the nozzle pointed towards the forehead. Do not angle the nozzle of the nasal spray towards the inner or outer wall of the nose. • Press the plunger firmly until the dose is administered and the spray is released into the nostril. • Note the time the Neffy device was administered. • Hand the used adrenaline device to the ambulance paramedics and advise them of the time of administration.
3.	Call an ambulance (000)
4.	If there is no improvement or severe symptoms progress (as described in the ASCIA Action Plan for Anaphylaxis (RED)), further adrenaline doses may be administered every five minutes, if other adrenaline autoinjectors are available.
5.	Contact the student's emergency contacts.
6.	The principal or a staff member allocated to do so must contact the Incident Support and Operations Centre (ISOC) on 1800 126 126 to report 'High' or Extreme' severity incidents to report the incident. Incidents assessed as 'Low' or 'Medium' can be reported directly into EduSafe Plus by the Principal or their allocated staff member.

If a student appears to be having a severe allergic reaction but has not been previously diagnosed with an allergy or being at risk of anaphylaxis, school staff should follow steps 2 – 5 as above.

For first time anaphylactic reactions, the school's general use adrenaline autoinjector device must be used. If the general use device is not immediately available in an anaphylaxis emergency, staff may use another student's adrenaline device, including the Epipen®, Anapen®, Jext® or Neffy® device. This may save a life. If another student's adrenaline device is used in an anaphylaxis emergency, the school will notify the parents of the student whose device was used and immediately replace the device.

Where possible, schools should consider using the correctly dosed adrenaline device depending on the weight of the student. However, in an emergency if there is no other option available, any device should be administered to the student.

Communication Plan

This policy will be available on the Frankston Heights Primary School website so that parents and other members of the school community can easily access information about the school's anaphylaxis management procedures.

The Principal is responsible for ensuring that all relevant staff, including casual relief staff, canteen staff and volunteers are aware of this policy and Frankston Heights Primary School procedures for anaphylaxis management. The policy will be communicated through staff induction and training materials, and twice yearly staff briefings. Casual relief staff and volunteers who are responsible for the care and/or supervision of students who are identified as being at risk of anaphylaxis will be notified via CRT form and/or verbally of their role in responding to an anaphylactic reaction and where required, the identity of students at risk.

The Principal is also responsible for ensuring relevant staff are trained and briefed in anaphylaxis management, consistent with the department's [Anaphylaxis Guidelines](#).

Staff training

Staff at Frankston Heights Primary School will receive appropriate training in anaphylaxis management, consistent with the Department's [Anaphylaxis Guidelines](#). This will include:

- First Aid Officer
- School staff who conduct classes attended by students who are at risk of anaphylaxis
- School staff who conduct specialist classes, admin staff, and any other member of school staff as required by the Principal.

Staff who are required to undertake training must:

- complete the ASCIA e-training course every 2 years
- have a school anaphylaxis supervisor verify their competency in the correct use of adrenaline devices (Epipen, Anapen, Jext and Neffy) in line with the Anaphylaxis Management School Training Checklist, within 30 days of completing this course
- have the ASCIA certificate signed by a school anaphylaxis supervisor.

Two staff members will be trained as school anaphylaxis supervisors to verify staff competency in the correct use of adrenaline devices (Epipen, Anapen, Jext and Neffy) and sign the ASCIA certificates. Trained staff must complete the following courses:

- the [ASCIA e-training course](#) every 2 years
- the Verifying the Correct Use of Adrenaline Injector Devices 22579VIC every 3 years (provided by Hero). Staff must also have a certificate in the [ASCIA e-training course](#), completed within the previous 12 months of enrolling into Verifying the Correct Use of Adrenaline Injector Devices 22579VIC
- complete the First Aid Management of Anaphylaxis 22578VIC every 3 years (provided by Hero).

Staff are also required to attend a briefing on anaphylaxis management and this policy at least twice per year (with the first briefing to be held at the beginning of the school year), facilitated by a staff member who has successfully completed an anaphylaxis management course within the last 2 years. Each briefing will address:

- the school's anaphylaxis management policy
- the causes, symptoms and treatment of anaphylaxis
- the identities of students diagnosed as being at risk of anaphylaxis, their allergens and the location of their Individual Anaphylaxis Management Plans and their medication/s
- discussion on staff anaphylaxis training and renewal

- how to use an adrenaline device, including hands-on practice with an adrenaline device trainer device (which does not contain adrenaline)
- the school's general first aid and emergency procedures
- the location of adrenaline devices prescribed for individual students that have been purchased by their family
- the location of adrenaline devices that the school has purchased for general use
- how to access on-going support and training.

When a new student enrolls at Frankston Heights Primary School who is at risk of anaphylaxis, the First Aid Officer will develop an interim plan in consultation with the student's parents and ensure that appropriate staff are trained and briefed as soon as possible.

A record of staff training courses and briefings will be maintained on the Frankston Heights Primary School OHS training planner.

The Principal will ensure that while students at risk of anaphylaxis are under the care or supervision of the school outside of normal class activities, including in the school yard, at camps and excursions, or at special event days, there is a sufficient number of school staff present who have been trained in anaphylaxis management.

COMMUNICATION

This policy will be communicated to our school community in the following ways:

- Available publicly on our school's website
- Staff induction processes and staff training
- Discussed at staff briefings/meetings as required
- School communication platform, Sentral

FURTHER INFORMATION AND RESOURCES

- The Department's Policy and Advisory Library (PAL):
 - [Anaphylaxis](#)
 - Allergies
 - First Aid for Students and Staff
 - Health Care Needs
 - Managing Reporting School Incidents (Including Emergencies)
 - Medication
- [Allergy & Anaphylaxis Australia](#)
- ASCIA Guidelines: [Schooling and childcare](#)
- [Hero HQ Anaphylaxis Management Training](#)
- Royal Children's Hospital: [Allergy and immunology](#)

RELATED FHPS POLICIES:

- Administration of Medication Policy
- First Aid Policy
- Health Care Needs Policy

POLICY REVIEW AND APPROVAL

The First Aid Officer and Assistant Principal under the direction of the Principal will complete the Department's Annual Risk Management Checklist for anaphylaxis management to assist with the evaluation and review of this policy and the support provided to students at risk of anaphylaxis.

Policy last reviewed	May 2026
Approved by	Principal
Next scheduled review date	May 2027